



Side Effects Guide

Your guide to understanding potential side effects associated with JEMPERLI*

*This information is not a substitute for your healthcare providers' advice. Be sure to speak to your doctor about your concerns and questions.

Not an actual patient taking JEMPERLI.

Please see Important Safety Information throughout, as well as full [Prescribing Information](#), including [Medication Guide](#), also available at JEMPERLI.com.

Jemperli 
(dostarlimab-gxly) Injection 500 mg



Table of Contents

You may experience side effects from JEMPERLI4

Potential side effects associated with JEMPERLI alone.....7

Potential side effects associated with JEMPERLI + carboplatin and paclitaxel (CP).....11

Approved Uses

JEMPERLI is a prescription medicine used to treat adults with:

- a kind of uterine cancer called endometrial cancer (EC)
 - JEMPERLI may be used in combination with the chemotherapy medicines, carboplatin and paclitaxel, and then after that JEMPERLI may be used alone:
 - when your cancer has spread outside your uterus (advanced) **or**,
 - your cancer has returned.
 - JEMPERLI may be used alone:
 - when a laboratory test shows that your tumor is mismatch repair deficient (dMMR), **and**
 - your cancer has returned, or it has spread (advanced EC), **and**
 - you have received chemotherapy that contains platinum and it did not work or is no longer working, **and**
 - your cancer cannot be treated by surgery or radiation.

It is not known if JEMPERLI is safe and effective in children.

Important Safety Information

JEMPERLI is a medicine that may treat certain cancers by working with your immune system. JEMPERLI can cause your immune system to attack normal organs and tissues in any area of your body and can affect the way they work. These problems can sometimes become severe or life-threatening and can lead to death. You can have more than one of these problems at the same time. These problems may happen anytime during treatment or even after your treatment has ended.

Please see additional Important Safety Information throughout, as well as full [Prescribing Information](#), including [Medication Guide](#), also available at JEMPERLI.com.

Jemperli 
(dostarlimab-gxly) Injection 500 mg



Not an actual patient taking JEMPERLI.

Important Safety Information (cont'd)

Call or see your healthcare provider (HCP) right away if you develop any new or worsening signs or symptoms, including:

Lung problems. Signs and symptoms may include cough, shortness of breath, or chest pain.

Intestinal problems. Signs and symptoms may include diarrhea or more bowel movements than usual; stools that are black, tarry, sticky, or have blood or mucus; or severe stomach-area (abdomen) pain or tenderness.

Liver problems. Signs and symptoms may include yellowing of your skin or the whites of your eyes, severe nausea or vomiting, pain on the right side of your stomach area (abdomen), dark urine (tea colored), or bleeding or bruising more easily than usual.

Please see additional Important Safety Information throughout, as well as full [Prescribing Information](#), including [Medication Guide](#), also available at JEMPERLI.com.

Jemperli 
(dostarlimab-gxly) Injection 500 mg

You may experience side effects from JEMPERLI

JEMPERLI is a medicine that may treat certain cancers by working with your immune system. JEMPERLI can cause your immune system to attack normal organs and tissues in any area of your body and can affect the way they work. These problems can sometimes become severe or life-threatening and can lead to death. You can have more than one of these problems at the same time. These problems may happen anytime during treatment or even after your treatment has ended.

Call or see your healthcare provider right away if you develop any new or worsening signs or symptoms, including:

Immune system problems

- Lung problems, like cough or shortness of breath
- Intestinal problems, like diarrhea or more bowel movements than usual, or severe stomach-area (abdomen) pain or tenderness
- Liver problems, like yellowing of your skin or the whites of your eyes
- Hormone gland problems, like headaches that will not go away or unusual headaches, or eye sensitivity to light
- Kidney problems, like swelling in your ankles or blood in your urine
- Skin problems, like rash or itching
- Problems in other organs and tissues, with symptoms including chest pain, persistent or severe muscle pain or weakness, or double vision

Other problems

- Infusion reactions that can sometimes be severe or life-threatening, with symptoms including chills or shaking, dizziness, or shortness of breath or wheezing
- Rejection of a transplanted organ
- Complications, including graft-versus-host-disease (GVHD), in people who have received a bone marrow (stem cell) transplant that uses donor stem cells (allogeneic)



These are not all of the signs and symptoms of problems that can happen with JEMPERLI. **Talk to your doctor to learn more.**

Please see additional Important Safety Information throughout, as well as full [Prescribing Information](#), including [Medication Guide](#), also available at JEMPERLI.com.

Jemperli 
(dostarlimab-gxly) Injection 500 mg

You may experience side effects from JEMPERLI (cont'd)

Getting medical treatment right away may help keep the problems on the previous page from becoming more serious.

- Your healthcare provider will check you for these problems during treatment with JEMPERLI
- If you have certain side effects, your healthcare provider may treat you with corticosteroid or hormone replacement medicines
- Your healthcare provider may also need to delay or completely stop treatment with JEMPERLI if you have severe side effects

In study participants receiving JEMPERLI + carboplatin and paclitaxel (CP):

- 19% stopped taking JEMPERLI completely due to side effects
- 37% had their treatment interrupted due to side effects

Of study participants who received JEMPERLI alone:

- 10% stopped taking JEMPERLI completely due to side effects
- 28% had their treatment interrupted due to side effects

Important Safety Information (cont'd)

Call or see your healthcare provider (HCP) right away if you develop any new or worsening signs or symptoms, including (cont'd):

Hormone gland problems. Signs and symptoms may include headaches that will not go away or unusual headaches, eye sensitivity to light, eye problems, rapid heartbeat, increased sweating, extreme tiredness, weight gain or weight loss, feeling more hungry or thirsty than usual, urinating more often than usual, hair loss, feeling cold, constipation, your voice gets deeper, dizziness or fainting, changes in mood or behavior such as decreased sex drive, irritability, or forgetfulness.

Kidney problems. Signs and symptoms may include change in the amount or color of your urine, blood in your urine, swelling in your ankles, or loss of appetite.

Skin problems. Signs and symptoms may include rash; itching; skin blistering or peeling; swollen lymph nodes; painful sores or ulcers in your mouth or in your nose, throat, or genital area; fever or flu-like symptoms.

Please see additional Important Safety Information throughout, as well as full [Prescribing Information](#), including [Medication Guide](#), also available at JEMPERLI.com.

Jemperli 
(dostarlimab-gxly) Injection 500 mg

You may experience side effects from JEMPERLI (cont'd)

The most common side effects of JEMPERLI + CP are nerve problems in your arms, hands, legs, and feet; tiredness; nausea; hair loss; joint pain; rash; constipation; diarrhea; stomach-area (abdomen) pain; shortness of breath; decreased appetite; urinary tract infections; vomiting.

The most common side effects of JEMPERLI when given alone include tiredness and weakness; low red blood cell count (anemia); diarrhea; nausea; constipation; vomiting.



Watch out for and talk to your doctor about

any and all side effects that you experience. They may be a sign of other more severe or life-threatening problems.

To learn more about side effects, visit [JEMPERLI.com](https://www.jemperli.com).

Important Safety Information (cont'd)

Problems can also happen in other organs and tissues. These are not all of the signs and symptoms of immune system problems that can happen with JEMPERLI. Call or see your HCP right away for any new or worse signs or symptoms. Signs and symptoms may include chest pain, irregular heartbeat, shortness of breath, swelling of ankles; confusion, sleepiness, memory problems, changes in mood or behavior, stiff neck, balance problems, tingling or numbness of the arms or legs; double vision, blurry vision, sensitivity to light, eye pain, changes in eyesight; persistent or severe muscle pain or weakness, muscle cramps; low red blood cells, bruising.

Infusion reactions that can sometimes be severe or life-threatening. Signs and symptoms of infusion reactions may include chills or shaking, itching or rash, flushing, shortness of breath or wheezing, dizziness, feel like passing out, fever, back or neck pain.

Please see additional Important Safety Information throughout, as well as full [Prescribing Information](#), including [Medication Guide](#), also available at [JEMPERLI.com](https://www.jemperli.com).

Jemperli 
(dostarlimab-gxly) Injection 500 mg

Potential side effects associated with JEMPERLI alone



Everyone's treatment experience is different. It's important to talk to your doctor when you first begin to experience side effects or if they worsen in any way. Please see below for more information on tips that your healthcare team may suggest for certain side effects, though these are not all the possible side effects of JEMPERLI. **Always let your healthcare provider know of any side effects you may experience and follow their guidance.**



Watch out for and talk to your doctor about any and all side effects that you experience. They may be a sign of other more severe or life-threatening problems. The side effects in this brochure are not all of the possible side effects you may experience with JEMPERLI. The tips listed in this brochure are not proven to help with these side effects.

Tiredness and Weakness

Talk to your care team about their recommendations for dealing with tiredness and weakness, which may include:

- Planning your day to prioritize things that are most important and spreading activities throughout the day
- Taking rest breaks or short naps rather than one long nap
- Planning an appropriate exercise program for you
- Aiming for 7 to 8 hours of sleep each night
- Avoiding things that interrupt sleep, such as caffeine and evening exercise

Important Safety Information (cont'd)

Rejection of a transplanted organ. Your HCP should tell you what signs and symptoms you should report and monitor you, depending on the type of organ transplant that you have had.

Complications, including graft-versus-host-disease (GVHD), in people who have received a bone marrow (stem cell) transplant that uses donor stem cells (allogeneic). These complications can be serious and can lead to death. These complications may happen if you underwent transplantation either before or after being treated with JEMPERLI. Your HCP will monitor you for these complications.

Please see additional Important Safety Information throughout, as well as full [Prescribing Information](#), including [Medication Guide](#), also available at JEMPERLI.com.

Jemperli 
(dostarlimab-gxly) Injection 500 mg

Potential side effects associated with JEMPERLI alone (cont'd)



Rely on your care team. They understand how you're feeling and have resources that may help.

Diarrhea

Talk to your care team about their recommendations for dealing with diarrhea, which may include:

- Avoiding fried foods and high-fiber foods like beans and whole grains
- Reducing the use of alcohol or tobacco
- Taking a medicine for diarrhea, which can be either prescription or over-the-counter and should be taken as instructed



Watch out for and talk to your doctor about any and all side effects that you experience. They may be a sign of other more severe or life-threatening problems. These are not all of the possible side effects you may experience with JEMPERLI. These tips are not proven to help with the side effects listed in this brochure.

Important Safety Information (cont'd)

Getting medical treatment right away may help keep these problems from becoming more serious. Your HCP will check you for these problems during treatment with JEMPERLI and may treat you with corticosteroid or hormone replacement medicines. If you have severe side effects, your HCP may also need to delay or completely stop treatment with JEMPERLI.

Please see additional Important Safety Information throughout, as well as full [Prescribing Information](#), including [Medication Guide](#), also available at JEMPERLI.com.

Jemperli 
(dostarlimab-gxly) Injection 500 mg



Not an actual patient taking JEMPERLI.

Potential side effects associated with JEMPERLI alone (cont'd)

Nausea

Talk to your care team about their recommendations for coping with nausea, which may include:

- Trying to avoid skipping meals or snacks—an empty stomach may make nausea worse
- Keeping dry toast, crackers, and other bland foods on hand
- Avoiding fatty, fried, spicy, or very sweet foods
- Sipping liquids slowly throughout the day
- Taking a medicine for nausea, which can be either prescription or over-the-counter and should be taken as instructed
- Trying aromatherapy, such as peppermint



Watch out for and talk to your doctor about any and all side effects that you experience. They may be a sign of other more severe or life-threatening problems. These are not all of the possible side effects you may experience with JEMPERLI. These tips are not proven to help with the side effects listed in this brochure.

Important Safety Information (cont'd)

Before receiving JEMPERLI, tell your HCP about all of your medical conditions, including immune system problems such as Crohn's disease, ulcerative colitis, or lupus; received an organ transplant; have received or plan to receive a stem cell transplant that uses donor stem cells (allogeneic); have received radiation treatment to your chest area; have a condition that affects your nervous system, such as myasthenia gravis or Guillain-Barré syndrome.

Please see additional Important Safety Information throughout, as well as full [Prescribing Information](#), including [Medication Guide](#), also available at JEMPERLI.com.

Jemperli 
(dostarlimab-gxly) Injection 500 mg

Potential side effects associated with JEMPERLI alone (cont'd)

Constipation

Talk to your care team about their recommendations for managing constipation, which may include:

- Trying to eat more high-fiber foods like fresh fruits and vegetables, oats, whole-grain bread, and brown rice
- Drinking plenty of fluids, but be sure to ask your care team how much is safe for you to drink
- Staying as active as you can with an appropriate exercise plan for you
- Taking a medicine for constipation, which can be either prescription or over-the-counter and should be taken as instructed

Vomiting

Talk to your care team about their recommendations for dealing with vomiting, which may include:

- Avoiding foods with strong or unpleasant smells
- Sipping liquids slowly until you feel good enough to drink more
- Talking with your care team about how to stay hydrated and how much is safe for you to drink



Watch out for and talk to your doctor about any and all side effects that you experience. They may be a sign of other more severe or life-threatening problems. These are not all of the possible side effects you may experience with JEMPERLI. These tips are not proven to help with the side effects listed in this brochure.

Important Safety Information (cont'd)

If you are pregnant or plan to become pregnant, tell your HCP. JEMPERLI can harm your unborn baby. If you are able to become pregnant, your HCP will give you a pregnancy test before you start treatment. Use an effective birth control method during treatment and for 4 months after your last dose of JEMPERLI. Tell your HCP right away if you become pregnant or think you may be pregnant during treatment with JEMPERLI.

If you are breastfeeding or plan to breastfeed, tell your HCP. It is not known if JEMPERLI passes into your breast milk. Do not breastfeed during treatment with JEMPERLI and for 4 months after your last dose.

Please see additional Important Safety Information throughout, as well as full [Prescribing Information](#), including [Medication Guide](#), also available at JEMPERLI.com.

Jemperli 
(dostarlimab-gxly) Injection 500 mg

Potential side effects associated with JEMPERLI + CP



Everyone's treatment experience is different. That's why it's important to talk to your doctor when you first begin to experience side effects or if they worsen in any way. Please see below for more information on tips that your healthcare team may suggest for certain side effects, though these are not all the possible side effects of JEMPERLI. **Always let your healthcare provider know of any side effects you may experience and follow their guidance.**

Nerve problems in your arms, hands, legs, and feet

Talk to your care team about their recommendations for dealing with nerve problems, which may include:

- Checking hands and feet regularly for any injuries, cuts, or sores
- Avoiding or limiting alcohol
- Performing gentle stretching and balancing exercises
- Protecting your hands and feet from extreme temperatures, like wearing shoes and warm clothes when it's cold and protecting your hands from heat while cooking
- Preventing falls by moving rugs out of your path and putting rails on the walls and in the bathroom
- Getting up slowly after sitting or lying down, especially if you feel dizzy
- Taking a medicine for pain, which may include a prescription or over-the-counter patch or cream, and should be taken as instructed



Watch out for and talk to your doctor about any and all side effects that you experience. They may be a sign of other more severe or life-threatening problems. The side effects in this brochure are not all of the possible side effects you may experience with JEMPERLI. The tips listed in this brochure are not proven to help with these side effects.

Important Safety Information (cont'd)

Tell your HCP about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Please see additional Important Safety Information throughout, as well as full [Prescribing Information](#), including [Medication Guide](#), also available at JEMPERLI.com.

Jemperli 
(dostarlimab-gxly) Injection 500 mg

Potential side effects associated with JEMPERLI + CP (cont'd)

Tiredness

Talk to your care team about their recommendations for dealing with tiredness, which may include:

- Planning your day to prioritize things that are most important and spreading activities throughout the day
- Taking rest breaks or short naps rather than one long nap
- Planning an appropriate exercise program for you
- Aiming for 7 to 8 hours of sleep each night
- Avoiding things that interrupt sleep, such as caffeine and evening exercise

Nausea

Talk to your care team about their recommendations for dealing with nausea, which may include:

- Trying to avoid skipping meals or snacks—an empty stomach may make nausea worse
- Keeping dry toast, crackers, and other bland foods on hand
- Avoiding fatty, fried, spicy, or very sweet foods
- Sipping liquids slowly throughout the day
- Taking a medicine for nausea, which can be either prescription or over-the-counter and should be taken as instructed
- Trying aromatherapy, such as peppermint



Watch out for and talk to your doctor about any and all side effects that you experience. They may be a sign of other more severe or life-threatening problems. These are not all of the possible side effects you may experience with JEMPERLI. These tips are not proven to help with the side effects listed in this brochure.

Important Safety Information (cont'd)

The most common side effects of JEMPERLI when given with carboplatin and paclitaxel include nerve problems in your arms, hands, legs, and feet; tiredness; nausea; hair loss; joint pain; rash; constipation; diarrhea; stomach-area (abdomen) pain; shortness of breath; decreased appetite; urinary tract infections; vomiting.

Please see additional Important Safety Information throughout, as well as full [Prescribing Information](#), including [Medication Guide](#), also available at JEMPERLI.com.

Jemperli 
(dostarlimab-gxly) Injection 500 mg

Potential side effects associated with JEMPERLI + CP (cont'd)

Hair loss

Talk to your care team about their recommendations for dealing with hair loss, which may include:

- Determining likelihood of hair loss and when it might begin
- Acknowledging that it's natural to feel upset about hair loss, and that some people choose to cut their hair short or shave their head before hair loss starts
- Trying wigs, scarves, and hats to see which option you prefer if it feels right for you

Joint pain

Talk to your care team about their recommendations for dealing with joint pain, which may include:

- Applying a cold compress, an ice pack, or a package of frozen vegetables to the painful area for 15 to 20 minutes a few times each day
- Applying a heating pad, soaking in a warm tub, or taking a warm shower to relax muscles and increase blood flow
- Visiting a physical therapist who can teach you gentle stretching exercises to ease muscle tension
- Taking a medicine for pain, which can be either prescription or over-the-counter and should be taken as instructed



Watch out for and talk to your doctor about any and all side effects that you experience. They may be a sign of other more severe or life-threatening problems. These are not all of the possible side effects you may experience with JEMPERLI. These tips are not proven to help with the side effects listed in this brochure.

Important Safety Information (cont'd)

The most common side effects of JEMPERLI when used alone include tiredness and weakness, low red blood cell count (anemia), diarrhea, nausea, constipation, vomiting.

These are not all of the possible side effects of JEMPERLI. Call your doctor for medical advice about side effects.

You are encouraged to report negative side effects of prescription drugs to the FDA by calling 1-800-FDA-1088 or visiting www.fda.gov/medwatch.

Please see additional Important Safety Information throughout, as well as full [Prescribing Information](#), including [Medication Guide](#), also available at JEMPERLI.com.

Jemperli 
(dostarlimab-gxly) Injection 500 mg



Potential side effects associated with JEMPERLI + CP (cont'd)

Rash

Talk to your care team about their recommendations for dealing with rash, which may include:

- Using mild lotions and moisturizers
- Using mild soap and a soft cloth to gently clean the area
- Protecting the area from heat and cold
- Staying out of the sun as much as you can
- Taking a medicine for rash, which can be either prescription or over-the-counter and should be taken as instructed



Watch out for and talk to your doctor about any and all side effects that you experience. They may be a sign of other more severe or life-threatening problems. These are not all of the possible side effects you may experience with JEMPERLI. These tips are not proven to help with the side effects listed in this brochure.

Important Safety Information

JEMPERLI is a medicine that may treat certain cancers by working with your immune system. JEMPERLI can cause your immune system to attack normal organs and tissues in any area of your body and can affect the way they work. These problems can sometimes become severe or life-threatening and can lead to death. You can have more than one of these problems at the same time. These problems may happen anytime during treatment or even after your treatment has ended.

Please see additional Important Safety Information throughout, as well as full [Prescribing Information](#), including [Medication Guide](#), also available at JEMPERLI.com.

Jemperli 
(dostarlimab-gxly) Injection 500 mg

Potential side effects associated with JEMPERLI + CP (cont'd)

Constipation

Talk to your care team about their recommendations for dealing with constipation, which may include:

- Trying to eat more high-fiber foods like fresh fruits and vegetables, oats, whole-grain bread, and brown rice
- Drinking plenty of fluids, but be sure to ask your care team how much is safe for you to drink
- Staying as active as you can with an appropriate exercise plan for you
- Taking a medicine for constipation, which can be either prescription or over-the-counter and should be taken as instructed

Diarrhea

Talk to your care team about their recommendations for dealing with diarrhea, which may include:

- Avoiding fried foods and high-fiber foods like beans and whole grains
- Reducing the use of alcohol or tobacco
- Taking a medicine for diarrhea, which can be either prescription or over-the-counter and should be taken as instructed



Watch out for and talk to your doctor about any and all side effects that you experience. They may be a sign of other more severe or life-threatening problems. These are not all of the possible side effects you may experience with JEMPERLI. These tips are not proven to help with the side effects listed in this brochure.

Important Safety Information (cont'd)

Call or see your healthcare provider (HCP) right away if you develop any new or worsening signs or symptoms, including:

Lung problems. Signs and symptoms may include cough, shortness of breath, or chest pain.

Intestinal problems. Signs and symptoms may include diarrhea or more bowel movements than usual; stools that are black, tarry, sticky, or have blood or mucus; or severe stomach-area (abdomen) pain or tenderness.

Please see additional Important Safety Information throughout, as well as full [Prescribing Information](#), including [Medication Guide](#), also available at JEMPERLI.com.

Jemperli 
(dostarlimab-gxly) Injection 500 mg

Potential side effects associated with JEMPERLI + CP (cont'd)

Decreased appetite

Talk to your care team about their recommendations for dealing with decreased appetite, which may include:

- Trying to eat small amounts more frequently, and taking advantage of times when you're hungry to eat more, if you feel full after eating a small amount
- Keeping snacks handy, so that you can eat when you're feeling up to it
- Asking a dietitian about shakes and instant drink mixes
- Scheduling your mealtimes, instead of relying on eating only when you feel hungry



Watch out for and talk to your doctor about any and all side effects that you experience. They may be a sign of other more severe or life-threatening problems. These are not all of the possible side effects you may experience with JEMPERLI. These tips are not proven to help with the side effects listed in this brochure.

Important Safety Information (cont'd)

Call or see your healthcare provider (HCP) right away if you develop any new or worsening signs or symptoms, including (cont'd):

Liver problems. Signs and symptoms may include yellowing of your skin or the whites of your eyes, severe nausea or vomiting, pain on the right side of your stomach area (abdomen), dark urine (tea colored), or bleeding or bruising more easily than usual.

Hormone gland problems. Signs and symptoms may include headaches that will not go away or unusual headaches, eye sensitivity to light, eye problems, rapid heartbeat, increased sweating, extreme tiredness, weight gain or weight loss, feeling more hungry or thirsty than usual, urinating more often than usual, hair loss, feeling cold, constipation, your voice gets deeper, dizziness or fainting, changes in mood or behavior such as decreased sex drive, irritability, or forgetfulness.

Kidney problems. Signs and symptoms may include change in the amount or color of your urine, blood in your urine, swelling in your ankles, or loss of appetite.

Please see additional Important Safety Information throughout, as well as full [Prescribing Information](#), including [Medication Guide](#), also available at JEMPERLI.com.

Jemperli 
(dostarlimab-gxly) Injection 500 mg

Potential side effects associated with JEMPERLI + CP (cont'd)

Urinary tract infection (UTI)

Talk to your care team about their recommendations for dealing with UTIs, which may include:

- Drinking plenty of liquids, so that urine is light yellow or clear, but be sure to ask your care team how much is safe for you to drink
- Preventing UTIs by going to the bathroom often, wearing cotton underwear and loose-fitting pants, and taking showers instead of baths
- Avoiding things that may make bladder problems worse, such as drinks with caffeine, alcohol, spicy foods, and tobacco products
- Taking a medication as recommended by your care team

Vomiting

Talk to your care team about their recommendations for dealing with vomiting, which may include:

- Avoiding foods with strong or unpleasant smells
- Talking with your care team about how to stay hydrated and how much is safe for you to drink
- Sipping liquids slowly until you feel good enough to drink more



Watch out for and talk to your doctor about any and all side effects that you experience. They may be a sign of other more severe or life-threatening problems. These are not all of the possible side effects you may experience with JEMPERLI. These tips are not proven to help with the side effects listed in this brochure.

Important Safety Information (cont'd)

Call or see your healthcare provider (HCP) right away if you develop any new or worsening signs or symptoms, including (cont'd):

Skin problems. Signs and symptoms may include rash; itching; skin blistering or peeling; swollen lymph nodes; painful sores or ulcers in your mouth or in your nose, throat, or genital area; fever or flu-like symptoms.

Please see additional Important Safety Information throughout, as well as full [Prescribing Information](#), including [Medication Guide](#), also available at JEMPERLI.com.

Jemperli 
(dostarlimab-gxly) Injection 500 mg



Want more resources that may help you on your treatment journey?

Visit [JEMPERLI.com](https://www.jemperli.com)

Not an actual patient taking JEMPERLI.

Important Safety Information (cont'd)

Problems can also happen in other organs and tissues. These are not all of the signs and symptoms of immune system problems that can happen with JEMPERLI. Call or see your HCP right away for any new or worse signs or symptoms. Signs and symptoms may include chest pain, irregular heartbeat, shortness of breath, swelling of ankles; confusion, sleepiness, memory problems, changes in mood or behavior, stiff neck, balance problems, tingling or numbness of the arms or legs; double vision, blurry vision, sensitivity to light, eye pain, changes in eyesight; persistent or severe muscle pain or weakness, muscle cramps; low red blood cells, bruising.

Infusion reactions that can sometimes be severe or life-threatening. Signs and symptoms of infusion reactions may include chills or shaking, itching or rash, flushing, shortness of breath or wheezing, dizziness, feel like passing out, fever, back or neck pain.

Rejection of a transplanted organ. Your HCP should tell you what signs and symptoms you should report and monitor you, depending on the type of organ transplant that you have had.

Please see additional Important Safety Information throughout, as well as full [Prescribing Information](#), including [Medication Guide](#), also available at [JEMPERLI.com](https://www.jemperli.com).

Trademarks are owned by or licensed to the GSK group of companies.

GSK

©2024 GSK or licensor.
PMUS-DSTBROC240019 October 2024
Produced in USA.

File Name: Side Effects Guide

Jemperli 
(dostarlimab-gxly) Injection 500 mg