

# BRIGHTER TOGETHER

## Test to know your dMMR status

Learn more about dMMR in endometrial cancer  
and a treatment that might be right for you >



dMMR, or mismatch repair deficient = an important biomarker status in endometrial cancer;  
endometrial cancer = cancer of the lining of the womb.

### APPROVED USE

JEMPERLI is a prescription medicine used to treat adults with a kind of uterine cancer called endometrial cancer and may be used when:

- your tumor has been shown by a laboratory test to be mismatch repair deficient (dMMR), **and**
- your cancer has returned, or it has spread (advanced cancer), **and**
- you have received chemotherapy that contains platinum and it did not work or is no longer working and your cancer cannot be treated by surgery or radiation.

It is not known if JEMPERLI is safe and effective in children.

### IMPORTANT SAFETY INFORMATION

JEMPERLI is a medicine that may treat certain cancers by working with your immune system. JEMPERLI can cause your immune system to attack normal organs and tissues in any area of your body and can affect the way they work. These problems can sometimes become severe or life-threatening and can lead to death. You can have more than one of these problems at the same time. These problems may happen anytime during treatment or even after your treatment has ended.

**Call or see your healthcare provider (HCP) right away if you develop any new or worsening of signs and symptoms, including:**

**Lung problems.** Signs and symptoms may include cough, shortness of breath, or chest pain.

Please see additional Important Safety Information throughout, as well as accompanying full Prescribing Information, also available at [JEMPERLI.com](http://JEMPERLI.com).

**Jemperli**   
(dostarlimab-gxly) injection 500 mg

# Understanding dMMR in endometrial cancer



- About **1 in 3 people** with endometrial cancer **have a biomarker called dMMR**, or mismatch repair deficient

**What is a biomarker?** It's a molecule found in the tissues or fluids of your body that shows if a condition, process, or disease is normal or abnormal



- **Knowing your dMMR status** can help your doctor:
  - Diagnose disease
  - Find a treatment that's right for you
  - Determine if there is an increased risk of cancer returning or spreading
  - Look for an increased risk of other cancers due to an inherited condition called Lynch syndrome



## IMPORTANT SAFETY INFORMATION (cont'd)

**Intestinal problems.** Signs and symptoms may include diarrhea or more bowel movements than usual; stools that are black, tarry, sticky, or have blood or mucus; or severe stomach-area (abdomen) pain or tenderness.

**Liver problems.** Signs and symptoms may include yellowing of your skin or the whites of your eyes, severe nausea or vomiting, pain on the right side of your stomach area (abdomen), dark urine (tea colored), or bleeding or bruising more easily than usual.

**Hormone gland problems.** Signs and symptoms may include headaches that will not go away or unusual headaches, eye sensitivity to light, eye problems, rapid heartbeat, increased sweating,

extreme tiredness, weight gain or weight loss, feeling more hungry or thirsty than usual, urinating more often than usual, hair loss, feeling cold, constipation, your voice gets deeper, dizziness or fainting, changes in mood or behavior such as decreased sex drive, irritability, or forgetfulness.

**Kidney problems.** Signs and symptoms may include change in the amount or color of your urine, blood in your urine, swelling in your ankles, or loss of appetite.

**Skin problems.** Signs and symptoms may include rash; itching; skin blistering or peeling; painful sores or ulcers in your mouth or in your nose, throat, or genital area; fever or flu-like symptoms; or swollen lymph nodes.



**Ask your doctor to test for dMMR**


Please see additional Important Safety Information throughout, as well as accompanying full Prescribing Information, also available at [JEMPERLI.com](http://JEMPERLI.com).

**Jemperli**   
(dostarlimab-gxly) Injection 500 mg

# How you may benefit from knowing your dMMR status

## If you have endometrial cancer with the dMMR biomarker, you might benefit from a treatment known as immunotherapy

- JEMPERLI is an immunotherapy option that may help your immune system fight cancer
  - JEMPERLI helps T cells, an important part of your immune system, find cancer cells so they can attack and fight the cancer



## IMPORTANT SAFETY INFORMATION (cont'd)

**Problems can also happen in other organs and tissues. These are not all of the signs and symptoms of immune system problems that can happen with JEMPERLI. Call or see your HCP right away for any new or worse signs or symptoms.**

Signs and symptoms may include chest pain, irregular heartbeat, shortness of breath, swelling of ankles; confusion, sleepiness, memory problems, changes in mood or behavior, stiff neck, balance problems, tingling or numbness of the arms or legs; double vision, blurry vision, sensitivity to light, eye pain, changes in eyesight; persistent or severe muscle pain or weakness, muscle cramps; low red blood cells, bruising.

**Infusion reactions that can sometimes be severe or life-threatening.** Signs and symptoms of infusion

reactions may include chills or shaking, itching or rash, flushing, shortness of breath or wheezing, dizziness, feel like passing out, fever, back or neck pain.

**Rejection of a transplanted organ.** Your HCP should tell you what signs and symptoms you should report and monitor you, depending on the type of organ transplant that you have had.

**Complications, including graft-versus-host-disease (GVHD), in people who have received a bone marrow (stem cell) transplant that uses donor stem cells (allogeneic).** These complications can be serious and can lead to death. These complications may happen if you underwent transplantation either before or after being treated with JEMPERLI. Your HCP will monitor you for these complications.



**Ask your doctor to test for dMMR and if JEMPERLI is right for you**

Please see additional Important Safety Information throughout, as well as accompanying full [Prescribing Information](#), also available at [JEMPERLI.com](#).

**Jemperli**   
(dostarlimab-gxly) Injection 500 mg

# What the experts are saying about dMMR testing

**National guidelines recommend testing for the dMMR biomarker if you have endometrial cancer**

## IMPORTANT SAFETY INFORMATION (cont'd)

### Getting medical treatment right away may help keep these problems from becoming more serious.

Your HCP will check you for these problems during treatment with JEMPERLI and may treat you with corticosteroid or hormone replacement medicines. If you have severe side effects, your HCP may also need to delay or completely stop treatment with JEMPERLI.

**Before receiving JEMPERLI, tell your HCP about all of your medical conditions, including** immune system problems such as Crohn's disease, ulcerative colitis, or lupus; received an organ transplant; have received or plan to receive a stem cell transplant that uses donor stem cells (allogeneic); have received radiation treatment to your chest area; have a condition that affects your nervous system, such as myasthenia gravis or Guillain-Barré syndrome.

**If you are pregnant or plan to become pregnant, tell your HCP.** JEMPERLI can harm your unborn baby. If you are able to become pregnant, your HCP will give you a pregnancy test before you start treatment. Use an effective birth control method during treatment and for 4 months after your last dose of JEMPERLI. Tell your HCP right away if you become pregnant or think you may be pregnant during treatment with JEMPERLI.

If you are breastfeeding or plan to breastfeed, tell your HCP. It is not known if JEMPERLI passes into your breast milk. Do not breastfeed during treatment with JEMPERLI and for 4 months after your last dose.

**Tell your HCP about all the medicines you take,** including prescription and over-the-counter medicines, vitamins, and herbal supplements.

**Common side effects of JEMPERLI** include tiredness and weakness, low red blood cell count (anemia), diarrhea, nausea, decreased number of certain white blood cells, decreased albumin in the blood, increase in certain liver blood tests, decreased salt (sodium) in the blood.

These are not all of the possible side effects of JEMPERLI. Call your doctor for medical advice about side effects.

You are encouraged to report negative side effects of prescription drugs to the FDA by calling 1-800-FDA-1088 or visiting [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

**Please see full Prescribing Information, including Medication Guide for patients.**



**Ask your doctor to test for dMMR and if JEMPERLI is right for you**

Trademarks are owned by or licensed to the GSK group of companies.



©2023 GSK or licensor.  
DSTLBN230001 March 2023  
Produced in USA. 0002-0022-42

**Jemperli**   
(dostarlimab-gxly) Injection 500 mg

Visit [JEMPERLI.com](http://JEMPERLI.com) for more information.